



TITLE:

# <Division of Biochemistry>Chemistry of Molecular Biocatalysts

AUTHOR(S):

---

CITATION:

<Division of Biochemistry>Chemistry of Molecular Biocatalysts. ICR  
Annual Report 2016, 23: 22-23

ISSUE DATE:

2016

URL:

<http://hdl.handle.net/2433/219059>

RIGHT:

Copyright © 2017 Institute for Chemical Research, Kyoto University

# Division of Biochemistry – Chemistry of Molecular Biocatalysts –

<http://www.scl.kyoto-u.ac.jp/~bunta/index-j.html>



Prof  
HIRATAKE, Jun  
(D Agr)



Assist Prof  
WATANABE, Bunta  
(D Agr)

## Students

ITO, Shota (M2)  
SAITO, Shingo (M2)

NAESHIRO, Kazuki (M2)  
LI, Xian (M2)


MAKINO, Katsunori (M1)

## Scope of Research

Life is the integration of numerous chemical interactions conducted by low molecular weight compounds and proteins. Our interest is understanding the mechanisms of these interactions from the viewpoint of organic chemistry. Our current research focuses on the following: a) Design and synthesis of a series of chemical probes in order to reveal not only ligand–protein and protein–protein interactions, but also behavior of low molecular weight compounds *per se*. b) Unraveling biosynthetic pathways of bioactive natural products at enzyme level. c) Application of our chemical probes in development of novel practical bioactive compounds.

### KEYWORDS

Enzyme Inhibitors  
 $\gamma$ -Glutamyl Transpeptidase  
Fructosyl Peptide Oxidase  
Diabetes Diagnosis  
Protein–Protein Interaction Inhibitors



## Selected Publications

Kamiyama, A.; Nakajima, M.; Han, L.; Wada, K.; Mizutani, M.; Tabuchi, Y.; Kojima-Yuasa, A.; Matsui-Yuasa, I.; Suzuki, H.; Fukuyama, K.; Watanabe, B.; Hiratake, J., Phosphonate-Based Irreversible Inhibitors of Human  $\gamma$ -Glutamyl Transpeptidase (GGT). GGsTop is a Non-Toxic and Highly Selective Inhibitor with Critical Electrostatic Interaction with an Active-Site Residue Lys562 for Enhanced Inhibitory Activity, *Bioorg. Med. Chem.*, **24**, 5340-5352 (2016).

Watanabe, B.; Minami, S.; Ishida, H.; Yoshioka, R.; Nakagawa, Y.; Morita, T.; Hayashi, K., Stereospecific Inhibitory Effects of CCG-1423 on the Cellular Events Mediated by Myocardin-Related Transcription Factor A, *PLOS ONE*, **10**, [e0136242-1]-[e0136242-16] (2015).

Watanabe, B.; Ichiiyanagi, A.; Hirokawa, K.; Gomi, K.; Nakatsu, T.; Kato, H.; Kajiyama, N., Synthesis and Inhibitory Activity of Substrate-Analog Fructosyl Peptide Oxidase Inhibitors, *Bioorg. Med. Chem. Lett.*, **25**, 3910-3913 (2015).

Nakajima, M.; Watanabe, B.; Han, L.; Shimizu, B.; Wada, K.; Fukuyama, K.; Suzuki, H.; Hiratake, J., Glutathione-Analogous Peptidyl Phosphorus Esters as Mechanism-Based Inhibitors of  $\gamma$ -Glutamyl Transpeptidase for Probing Cysteiny-Glycine Binding Site, *Bioorg. Med. Chem.*, **22**, 1176-1194 (2014).


Hayashi, K.; Watanabe, B.; Nakagawa, Y.; Minami, S.; Morita, T., RPEL Proteins are the Molecular Targets for CCG-1423, an Inhibitor of Rho Signaling, *PLOS ONE*, **9**, [e89016-1]-[e89016-12] (2014).

## Determination of a Key Residue of $\gamma$ -Glutamyl Transpeptidase for Substrate Recognition


$\gamma$ -Glutamyl transpeptidase (GGT) plays a central role in homeostasis of antioxidant tripeptide glutathione, and has been implicated in a vast array of physiological disorders. In this study, we synthesized a series of mechanism-based GGT inhibitors to probe electrostatic interactions between the acceptor site residues of GGT and substrates. Our chemical, enzymological, and molecular biological approaches revealed that 3-hydroxyphenylacetic acid is an excellent mimic of the cysteinylglycine moiety of glutathione, and Lys562 of human GGT strongly recognizes their negative charge on the carboxy group (Figure 1). We demonstrated that this interaction considerably enhances the human GGT specificity of our inhibitor named GGsTop. GGsTop exhibited no inhibitory activity at 10 mM on a representative member of glutamine-dependent amidotransferases essential for a wide range of biosynthetic pathway, and showed no cytotoxicity toward human fibroblasts and hepatic stellate cells up to 1 mM.

## Substrate-Analog Fructosyl Peptide Oxidase Inhibitors

Fructosyl peptide oxidase (FPOX) is widely used in the



**Figure 1.** Proposed binding mode of GGsTop, glutathione, and acceptor substrate (Gly-Gly) to Lys562.




**Figure 2.** Chemical structures and inhibitory activity of FPOX inhibitors.

area of diabetes diagnosis today. In this study, we designed and synthesized its substrate-analog inhibitors in order to unveil the substrate recognition mechanism of FPOX by X-ray diffraction analysis of enzyme-inhibitor co-crystals. Kinetic study revealed that our substrate analogs act as competitive inhibitors with  $K_i$  values ranging from 11.1 to 782  $\mu$ M (Figure 2). Co-crystallization of the enzyme with our inhibitors in order to determine the three-dimensional structure of FPOX is now in progress.

## Molecular Mechanism of Myocardin-Related Transcription Factor A Inhibitors

Myocardin-related transcription factor A (MRTF-A) plays a pivotal role in epidermal-mesenchymal transition. Inhibition of its nuclear transport is regarded as one of the attractive therapeutic targets since MRTF-A is closely associated with cancer and tissue fibrosis. In this study, we revealed that CCG-1423, originally developed as a Rho inhibitor, binds to the nuclear localization signal of MRTF-A and inhibits its nuclear transport mediated by importin- $\alpha$ / $\beta$ 1 (Figure 3). We also demonstrated that CCG-1423 inhibits a migration of melanoma cells triggered by MRTF-A activation, and the potency is affected by the stereochemistry of CCG-1423. The difference is elucidated by the binding manner of each stereoisomer to MRTF-A that speculated by a molecular modeling approach.



**Figure 3.** Molecular mechanism of MRTF-A inhibitor CCG-1423.